



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Matrix multiplication is determined by orthogonality and trace

Citation for published version:

Heunen, C & Horsman, C 2013, 'Matrix multiplication is determined by orthogonality and trace', *Linear algebra and its applications*, vol. 439, no. 12, pp. 4130-4134. <https://doi.org/10.1016/j.laa.2013.09.039>

Digital Object Identifier (DOI):

[10.1016/j.laa.2013.09.039](https://doi.org/10.1016/j.laa.2013.09.039)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Linear algebra and its applications

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



MATRIX MULTIPLICATION IS DETERMINED BY ORTHOGONALITY AND TRACE

CHRIS HEUNEN AND CLARE HORSMAN

ABSTRACT. Any associative bilinear multiplication on the set of n -by- n matrices over some field of characteristic not two, that makes the same vectors orthogonal and has the same trace as ordinary matrix multiplication, must be ordinary matrix multiplication or its opposite.

1. INTRODUCTION

Matrix multiplication is the fundamental operation of linear algebra, modelling composition of linear maps in terms of coordinates. This article characterises matrix multiplication simply in terms of orthogonality and trace, in the following way. Suppose there were another way to combine two matrices $x, y \in \mathbb{M}_n$ into a new one $x \star y \in \mathbb{M}_n$, that resembles composition of linear maps in the sense that it is associative, bilinear, and respects the identity matrix:

- (A) “*Associativity*”: $x \star (y \star z) = (x \star y) \star z$ for all $x, y, z \in \mathbb{M}_n$;
- (B) “*Bilinearity*”: $(\lambda x) \star y = \lambda(x \star y) = x \star (\lambda y)$, $(x + y) \star z = (x \star z) + (y \star z)$, and $x \star (y + z) = (x \star y) + (x \star z)$ for scalars λ and $x, y, z \in \mathbb{M}_n$;
- (I) “*Identity*”: $x \star 1 = x$ for all $x \in \mathbb{M}_n$, where $1 \in \mathbb{M}_n$ is the identity matrix.

To these basic requirements we add two properties that, at first sight, do not fix much information about $x \star y$. First, we require $x \star y$ to have the same trace as ordinary matrix multiplication xy :

- (T) “*Trace*”: $\text{Tr}(x \star y) = \text{Tr}(xy)$ for all $x, y \in \mathbb{M}_n$.

Second, we require that \star makes the same vectors orthogonal as ordinary matrix multiplication. Formulated algebraically:

- (O) “*Orthogonality*”: $x \star y = 0$ when $xy = yx = 0$, $xx = x$, and $yy = y$, for $x, y \in \mathbb{M}_n$ that have rank one.

These assumptions already imply that $x \star y$ must equal the ordinary matrix multiplication, or its opposite, after all. That is, we will prove the following theorem, for any scalar field \mathbb{F} of characteristic not two.

Theorem 1. *A function $\star: \mathbb{M}_n \times \mathbb{M}_n \rightarrow \mathbb{M}_n$ satisfies (A), (B), (I), (T) and (O) if and only if either $x \star y = xy$ for all $x, y \in \mathbb{M}_n$, or $x \star y = yx$ for all $x, y \in \mathbb{M}_n$.*

This fits in the recent programme of results surrounding linear preserver problems, *i.e.* linear maps that preserve zero products, commutativity, etc. [4, 6, 1, 7]. Indeed, we will rely on one of those results [1]. Our original motivation came from quantum theory, where the above properties arise as desiderata for a possible noncommutative extension of Bayesian inference [5, 3].

2010 *Mathematics Subject Classification.* 15A03, 15A04, 15A99, 15A86.

Key words and phrases. Matrix multiplication, orthogonality, trace, zero products.

2. THE MAIN RESULT

Proposition 2. *A map $\star: \mathbb{M}_n \times \mathbb{M}_n \rightarrow \mathbb{M}_n$ meets (B), (I), and (O) if and only if*

$$(1) \quad x \star y = xy + g(xy - yx)$$

for some linear map $g: \mathbb{M}_n \rightarrow \mathbb{M}_n$ and all $x, y \in \mathbb{M}_n$.

Proof. Assume (B), (I) and (O). Then [1, Theorem 2.2] applies, giving linear maps $f, g: \mathbb{M}_n \rightarrow \mathbb{M}_n$ with $x \star y = f(xy) + g(yx)$. Furthermore, $x = x \star 1 = f(x1) + g(1x)$, and so $f(x) = x - g(x)$. Therefore $x \star y = f(xy) + g(yx) = xy + g(yx - xy)$. Taking the negative of g now gives (1).

If \star is of the form (1), then it is easy to show that (B), (I), and (O) hold. \square

Lemma 3. *Suppose \star satisfies (B), (I), and (O). Then (T) holds if and only if g sends traceless matrices to traceless matrices.*

Proof. Property (T) is equivalent to $\text{Tr}(g(xy - yx)) = 0$ for all $x, y \in \mathbb{M}_n$. But matrices of the form $xy - yx$ are precisely those with trace zero [2]. \square

Write e_{ij} for the standard matrix units, so that $e_{ij}e_{kl} = \delta_{jk}e_{il}$. Because of bilinearity, property (A) holds precisely when it holds for $x = e_{ab}$, $y = e_{cd}$, $z = e_{ef}$ for all $a, b, c, d, e, f \in \{1, \dots, n\}$. Via Proposition 2, property (A) comes down to

$$(2) \quad \begin{aligned} & -\delta_{de}\delta_{af}g(e_{cb}) + \delta_{de}e_{ab} \star g(e_{cf}) - \delta_{cf}e_{ab} \star g(e_{ed}) \\ & = -\delta_{af}\delta_{bc}g(e_{ed}) + \delta_{bc}g(e_{ad}) \star e_{ef} - \delta_{ad}g(e_{cb}) \star e_{ef} \end{aligned}$$

for all $a, b, c, d, e, f \in \{1, \dots, n\}$. We will use this formula very often below.

By linearity of g , we may write $g(e_{ij}) = \sum_{k,l=1}^n G_{kl,ij}e_{lk}$ for entries $G_{kl,ij} \in \mathbb{F}$. By convention, we will write $g(ij)_{kl}$ for $G_{kl,ij}$, and $g(ii - jj)_{kl}$ for $G_{kl,ii} - G_{kl,jj}$.

Lemma 4. *For distinct $i, j, k, l \in \{1, \dots, n\}$:*

$$(3) \quad g(ij)_{kl} = 0,$$

$$(4) \quad g(ii - jj)_{kl} = 0.$$

For distinct $i, j, k \in \{1, \dots, n\}$:

$$(5) \quad g(ij)_{jk} = 0,$$

$$(6) \quad g(ij)_{kj} = 0,$$

$$(7) \quad g(ij)_{ik} = 0,$$

$$(8) \quad g(ij)_{ki} = 0,$$

$$(9) \quad g(ij)_{kk} = 0,$$

$$(10) \quad g(ii - jj)_{kk} = 0,$$

$$(11) \quad g(ii - jj)_{ik} = 0,$$

$$(12) \quad g(ii - jj)_{ki} = 0,$$

$$(13) \quad g(ii - jj)_{kj} = 0,$$

$$(14) \quad g(ii - jj)_{jk} = 0.$$

For distinct $i, j \in \{1, \dots, n\}$:

$$(15) \quad g(ij)_{jj} = 0,$$

$$(16) \quad g(ij)_{ii} = 0,$$

$$(17) \quad g(ij)_{ij} = 0,$$

$$(18) \quad g(ii - jj)_{ij} = 0,$$

$$(19) \quad g(ii - jj)_{ji} = 0.$$

Proof. All these equations are derived following the same pattern. Let i, j, k, l be distinct. Taking $a = k, b = l, c = i$, and $d = e = f = j$ in (2) results in $e_{kl} \star g(e_{ij}) = 0$. Taking the trace of both sides and using property (T) now shows $g(ij)_{kl} = \text{Tr}(g(e_{ij})e_{kl}) = \text{Tr}(e_{kl}g(e_{ij})) = \text{Tr}(e_{kl} \star g(e_{ij})) = 0$, establishing (3).

Similarly, taking $a = k, b = l, c = f = i$, and $d = e = j$ in (2) results in $e_{kl} \star g(e_{ii} - e_{jj}) = 0$, from which (4) follows by taking the trace of both sides.

For (5)–(19) we simply list the appropriate choice of indices:

- (5): $c = d = e = i, a = f = j, b = k$;
- (6): $c = d = e = i, b = f = j, a = k$;
- (7): $a = c = i, d = e = f = j, b = k$;
- (8): $a = f = i, d = j, b = c = e = k$;
- (9): $c = d = e = i, f = j, a = b = k$;
- (10): $c = f = i, d = e = j, a = b = k$;
- (11): $a = c = f = i, d = e = j, b = k$;
- (12): $b = c = f = i, d = e = j, a = k$; use (5);
- (13): $c = f = i, b = d = e = j, a = k$;
- (14): $c = f = i, a = d = e = j, b = k$; use (8);
- (15): $c = d = e = i, a = b = f = j$;
- (16): $a = b = c = i, d = e = f = j$; use (15);
- (17): $a = c = i, b = d = e = f = j$;
- (18): $a = c = f = i, b = d = e = j$;
- (19): $b = c = f = i, a = d = e = j$. □

Lemma 5. *There is $\lambda \in \{0, -1\}$ with $g(ii - jj)_{ii} = -g(ii - jj)_{jj} = g(ij)_{ji} = \lambda$ for distinct $i, j \in \{1, \dots, n\}$. Moreover, there is $z \in \mathbb{M}_n$ with $g(ii) = \lambda e_{ii} + z$ for all i .*

Proof. For any $k, l \in \{1, \dots, n\}$, the entry $g(ij)_{kl}$ can only be nonzero when $k = j$ and $l = i$ by (3), (5)–(9) and (15)–(17). Let $\lambda_{ij} \in \mathbb{F}$ be that entry: $g(ij) = \lambda_{ij}e_{ij}$.

Taking $b = c = e = f = i$ and $a = d = j$ in (2) leads to $g(ii - jj)_{ii} = g(ij)_{ji}$. Similarly, $a = d = i$ and $b = c = e = f = j$ lead to $g(jj - ii)_{jj} = g(ji)_{ij}$. Next, $b = c = d = e = i$ and $a = f = j$ show that $g(ij)_{ji} = g(ji)_{ij}$. Therefore

$$g(ii - jj)_{ii} = g(ij)_{ji} = \lambda_{ij}, \quad g(ii - jj)_{jj} = -g(ij)_{ji} = -\lambda_{ij}.$$

Combining this with (4), (10)–(14), (18) and (19) shows $g(ii - jj) = \lambda_{ij}(e_{ii} - e_{jj})$.

We may write $g(ii) = \lambda_i e_{ii} + z_i$ for $\lambda_i \in \mathbb{F}$, and $z_i \in \mathbb{M}_n$ linearly independent from e_{ii} . Then $\lambda_{ij}(e_{ii} - e_{jj}) = g(ii - jj) = \lambda_i e_{ii} - \lambda_j e_{jj} + (z_i - z_j)$. It follows that z_i does not depend on i , and we simply write z instead. Similarly, $\lambda_{ij} = \lambda_i = \lambda_j$ does not depend on i or j , and we may simply write λ for λ_{ij} . Hence $g(ii) = \lambda e_{ii} + z$.

It follows from the choice of indices for (18) that $g(ij) = e_{ij} \star g(ii - jj)$, and so

$$g(ij) = \lambda e_{ij} \star (e_{ii} - e_{jj}) = \lambda(-g(ij) - e_{ij} - g(ij)) = -2\lambda g(ij) - \lambda e_{ij}.$$

Hence $(1 + 2\lambda)g(ij) = -\lambda e_{ij}$. But $g(ij) = \lambda e_{ij}$ by definition, so $\lambda \in \{0, -1\}$. □

Proof of Theorem 1. Lemma 5 gives $\lambda \in \{0, -1\}$ and $z \in \mathbb{M}_n$ with $g(ii) = \lambda e_{ii} + z$. Let $x \in \mathbb{M}_n$, say $x = \sum_{i,j=1}^n \chi_{ij} e_{ij}$ for $\chi_{ij} \in \mathbb{F}$. It follows from Lemma 4 that

$$\begin{aligned} g(x) &= \sum_{i,j=1}^n \chi_{ij} g(ij) = \sum_{i=1}^n \chi_{ii} g(ii) + \sum_{i \neq j} \chi_{ij} g(ij) \\ &= \sum_{i=1}^n (\lambda \chi_{ii} e_{ii} + \chi_{ii} z) + \sum_{i \neq j} \lambda \chi_{ij} e_{ij} \\ &= \lambda \sum_{i,j=1}^n \chi_{ij} e_{ij} + \sum_{i=1}^n \chi_{ii} z \\ &= \lambda x + \text{Tr}(x)z. \end{aligned}$$

Therefore $x \star y = xy + \lambda(xy - yx)$. \square

Remark 6. The restriction of the base field \mathbb{F} to characteristic not two is necessary. For example, if $\mathbb{F} = \mathbb{Z}_2$, then

$$(20) \quad \begin{pmatrix} \alpha & \beta \\ \gamma & \varepsilon \end{pmatrix} \star \begin{pmatrix} \zeta & \eta \\ \theta & \iota \end{pmatrix} = \begin{pmatrix} \alpha\zeta + \gamma\eta & \beta\zeta + \varepsilon\eta \\ \alpha\theta + \gamma\iota & \beta\theta + \varepsilon\iota \end{pmatrix}$$

satisfies (A), (B), (I), (O), and (T), but differs from matrix multiplication or its opposite. In fact, apart from $g = 0, 1$, this is the only one of the form (1), for *e.g.*

$$g \begin{pmatrix} \alpha & \beta \\ \gamma & \varepsilon \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \alpha \end{pmatrix}.$$

ACKNOWLEDGEMENTS

The physical considerations leading to Theorem 1 were originally raised by Jonathan Barrett and Matthew Pusey, and are discussed in [3]. Chris Heunen was supported by the Office of Naval Research under grant N000141010357. Clare Horsman was supported by the CHIST-ERA DIQIP project, and by the FQXi Large Grant “Time and the Structure of Quantum Theory”.

REFERENCES

- [1] J. Alaminos, M. Brešar, J. Extremera, and A. R. Villena. On bilinear maps determined by rank one idempotents. *Lin. Alg. Appl.*, 432:738–743, 2010.
- [2] A. A. Albert and B. Muckenhoupt. On matrices of trace zero. *Mich. Math. J.*, 4(1):1–3, 1957.
- [3] J. Barrett, C. Heunen, C. Horsman, M. Pusey, and R. W. Spekkens. Causal inference in quantum mechanics. *forthcoming*, 2013.
- [4] M. Brešar and P. Šemrl. On bilinear maps on matrices with applications to commutativity preservers. *J. Alg.*, 301:803–837, 2006.
- [5] M. S. Leifer and R. W. Spekkens. A Bayesian approach to compatibility, improvement, and pooling of quantum states. *arXiv:1110.1085*, 2011.
- [6] L. Molnár. *Selected Preserver Problems on Algebraic Structures of Linear Operators and on Function Spaces*. Springer, 2007.
- [7] D. Wang, X. Li, and H. Ge. Idempotent elements determined matrix algebras. *Lin. Alg. Appl.*, 435:2889–2895, 2011.

DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF OXFORD
E-mail address: {chris.heunen, clare.horsman}@cs.ox.ac.uk